

ApoB Rabbit mAb

货号: B31179

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IP FC
推荐浓度	WB: 1:500 - 1:2000 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	515kDa
实测分子量	515kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse plasma,Rat plasma
细胞定位	Cytoplasm,Secreted
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human ApoB.
序列	LHFSNVFRSVMAPFTMTIDAHTNGNGKLALWGEHTGQLYSKFLKAEPLAFTFSHDYKGSTSHHLVSRKSISAALCHKVSA LLTPAEQTGTWKLKTQFNNNEYSQDLDAYNTKDKIGVELTGRTLADLTLLDSPIKVPLLLSEPINIIDALEMRDAVEKPQEFTI VAFVKYDKNQDVHSINLPFFETLQEYFERNRQTIIVVLENVQRNLKHINIDQFVRKYRAALGKLPQQANDYLNFSNWERQV SHAKEKLTALTKKYRITENDIQIALDDAKINFNEKLSQLQTYMIQFDQYIKDSYD

靶点信息

研究背景	This gene product is the main apolipoprotein of chylomicrons and low density lipoproteins. It occurs in plasma as two main isoforms, apoB-48 and apoB-100: the former is synthesized exclusively in the gut and the latter in the liver. The intestinal and the hepatic forms of apoB are encoded by a single gene from a single, very long mRNA. The two isoforms share a common N-terminal sequence. The shorter apoB-48 protein is produced after RNA editing of the apoB-100 transcript at residue 2180 (CAA->UAA), resulting in the creation of a stop codon, and early translation termination. Mutations in this gene or its regulatory region cause hypobetalipoproteinemia, normotriglyceridemic hypobetalipoproteinemia, and hypercholesterolemia due to ligand-defective apoB, diseases affecting plasma cholesterol and apoB levels.
基因ID	338
基因名	APOB
Swiss	P04114
别名	APOB;FLDB;LDLCQ4;apoB-100;apoB-48

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn