

ATP1B1 Rabbit mAb

货号: B31205

产品信息

| 反应 | Human,Mouse,Rat |
|-------|--|
| | папап, новернае |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IHC |
| 推荐浓度 | WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 |
| 理论分子量 | 40-50kDa |
| 实测分子量 | 40-50kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | U2OS,Mouse liver,Mouse brain,Rat brain |
| 细胞定位 | Cell membrane,Single-pass type II membrane protein |
| 纯化 | Affinity purification |


抗原信息

| 抗原信息 | Recombinant fusion protein corresponding to Human ATP1B1. |
|------|---|
| 序列 | YNPNVLPVQCTGKRDEDKDKVGNVEYFGLGNSPGFPLQYYPYYGKLLQPKYLQPLLAVQFTNLTMDTEIRIECKAYGENIG YSEKDRFQGRFDVKIEVKS |

靶点信息

| 研究背景 | The protein encoded by this gene belongs to the family of Na+/K+ and H+/K+ ATPases beta chain protein s, and to the subfamily of Na+/K+ -ATPases. Na+/K+ -ATPase is an integral membrane protein responsible for establishing and maintaining the electrochemical gradients of Na and K ions across the plasma membrane. These gradients are essential for osmoregulation, for sodium-coupled transport of a variety of organic and inorganic molecules, and for electrical excitability of nerve and muscle. This enzyme is composed of two subunits, a large catalytic subunit (alpha) and a smaller glycoprotein subunit (beta). The beta subunit regulates, through assembly of alpha/beta heterodimers, the number of sodium pumps transported to the plasma membrane. The glycoprotein subunit of Na+/K+ -ATPase is encoded by multiple genes. This gene encodes a beta 1 subunit. Alternatively spliced transcript variants encoding different isoforms have been described, but their biological validity is not known. [provided by RefSeq, Mar 2010] |
|--------------|--|
| 基因 ID | 481 |
| 基因名 | ATP1B1 |
| Swiss | P05026 |
| 别名 | ATP1B |

产品验证



Western blot analysis of ATP1B1 expressed in Mouse liver,Rat brain,U2OS using ATP1B1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lan e. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn