

RNF7 Rabbit mAb

货号: B31228

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	9kDa/10kDa/11kDa/12kDa
实测分子量	13kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	human liver tissue, human heart tissue, human skeletal muscle tissue, human testis tissue
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human RNF7.
序列	MADVEDGEETCALASHSGSSGSKSGGDKMFSLKKWNAVAMWSWDVECDTCAICRVQVMDACLRCQAENKQEDCVV VWGECNHSFHNCCMSLWVKQNNRCPLCQQDWVVQRIGK

靶点信息

研究背景	The protein encoded by this gene is a highly conserved ring finger protein. It is an essential subunit of SK P1-cullin/CDC53-F box protein ubiquitin ligases, which are a part of the protein degradation machinery im portant for cell cycle progression and signal transduction. This protein interacts with, and is a substrate of , casein kinase II (CSNK2A1/CKII). The phosphorylation of this protein by CSNK2A1 has been shown to pro
	mote the degradation of IkappaBalpha (CHUK/IKK-alpha/IKBKA) and p27Kip1(CDKN1B). Alternatively splic ed transcript variants encoding distinct isoforms have been reported.

基因 ID	9616
基因名	RNF7
Swiss	Q9UBF6
别名	RNF7;CKBBP1;ROC2;SAG

产品验证



Western blot analysis of RNF7 expressed in human liver tissue,human heart tissue,human skeletal muscl e tissue using RNF7 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:50 00. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanc ed Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn