

# Alpha Synuclein Rabbit mAb

货号: B31234

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	11kDa/13kDa/14kDa
实测分子量	20kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,Mouse brain,Rat brain
细胞定位	Cell junction,Cytoplasm,Membrane,Nucleus,Secreted,cytosol,synapse
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Alpha Synuclein.
序列	TKEQVTNVGGAVVTGVTAVAQKTVEGAGSIAAATGFVKKDQLGKNEEGAPQEGILEDMPVDPDNEAYEMPSEEGYQDYE PEA

靶点信息

研究背景	Alpha-synuclein is a member of the synuclein family, which also includes beta- and gamma-synuclein. Sy nucleins are abundantly expressed in the brain and alpha- and beta-synuclein inhibit phospholipase D2 s electively. SNCA may serve to integrate presynaptic signaling and membrane trafficking. Defects in SNCA have been implicated in the pathogenesis of Parkinson disease. SNCA peptides are a major component of amyloid plaques in the brains of patients with Alzheimer's disease. Alternatively spliced transcripts encod ing different isoforms have been identified for this gene.
基因ID	6622
基因名	SNCA
Swiss	P37840
别名	PD1;NACP;PARK1;PARK4;SNCA

## 产品验证



Western blot analysis of Alpha Synuclein expressed in 293T,Mouse brain,Rat brain using Alpha Synuc lein Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: www.ablybio.cn