

# Ret Rabbit mAb

货号: B31274

### 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	119kDa/124kDa
实测分子量	150,175kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	K-562
细胞定位	Cell membrane,Endosome membrane,Single-pass type I membrane protein
纯化	Affinity purification


### 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Ret.
序列	DSYVK

靶点信息

研究背景	This gene, a member of the cadherin superfamily, encodes one of the receptor tyrosine kinases, which ar e cell-surface molecules that transduce signals for cell growth and differentiation. This gene plays a cruci al role in neural crest development, and it can undergo oncogenic activation in vivo and in vitro by cytoge netic rearrangement. Mutations in this gene are associated with the disorders multiple endocrine neoplasi a, type IIA, multiple endocrine neoplasia, type IIB, Hirschsprung disease, and medullary thyroid carcinoma . Two transcript variants encoding different isoforms have been found for this gene. Additional transcript variants have been described but their biological validity has not been confirmed.
基因 <b>ID</b>	5979
基因名	RET
Swiss	P07949
别名	RET;CDHF12;CDHR16;HSCR1;MEN2A;MEN2B;MTC1;PTC;RET-ELE1;RET51

### 产品验证



Western blot analysis of Ret expressed in K-562 using Ret Rabbit mAb at 1:1000. Secondary antibody: HR P Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: www.ablybio.cn