

CD24 Rabbit mAb

货号: B31288

产品信息

| | |
|-------|---|
| 反应 | Human |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IP |
| 推荐浓度 | WB: 1:500 - 1:2000 IP: 1:20 - 1:50 |
| 理论分子量 | 8kDa |
| 实测分子量 | 45kDa/70kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | B cells,MCF7,SKOV3,Jurkat,Raji,Mouse brain |
| 细胞定位 | Cell membrane,GPI-anchor,Lipid-anchor |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|--|
| 抗原信息 | Recombinant fusion protein corresponding to Human CD24. |
| 序列 | MGRAMVARLGLGLLLLALLLPTQIYSSETTTGTSSNSSQSTSNSGLAPNPTNATTKAAGGALQSTASLFVVSLSLLHLYS |

靶点信息

| | |
|------|--|
| 研究背景 | This gene encodes a sialoglycoprotein that is expressed on mature granulocytes and B cells and modulates growth and differentiation signals to these cells. The precursor protein is cleaved to a short 32 amino acid mature peptide which is anchored via a glycosyl phosphatidylinositol (GPI) link to the cell surface. This gene was missing from previous genome assemblies, but is properly located on chromosome 6. Non-transcribed pseudogenes have been designated on chromosomes 1, 15, 20, and Y. Alternative splicing results in multiple transcript variants. |
| 基因ID | 100133941 |

| | |
|-------|--------------------------------|
| 基因名 | CD24 |
| Swiss | P25063 |
| 别名 | CD24;CD24A;CD24 molecule;CD247 |

产品验证



Western blot analysis of CD24 expressed in B cells,MCF7,SKOV3 using CD24 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn