

CD68 Rabbit mAb

货号: B31317

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	31kDa/34kDa/37kDa
实测分子量	37kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,Raji,THP-1,RAW264.7,Mouse liver
细胞定位	lysosome,plasma membrane
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein corresponding to Human CD68.
序列	


靶点信息

研究背景

This gene encodes a 110-kD transmembrane glycoprotein that is highly expressed by human monocytes and tissue macrophages. It is a member of the lysosomal/endosomal-associated membrane glycoprotein (LAMP) family. The protein primarily localizes to lysosomes and endosomes with a smaller fraction circula ting to the cell surface. It is a type I integral membrane protein with a heavily glycosylated extracellular d omain and binds to tissue- and organ-specific lectins or selectins. The protein is also a member of the sca venger receptor family. Scavenger receptors typically function to clear cellular debris, promote phagocyt osis, and mediate the recruitment and activation of macrophages. Alternative splicing results in multiple t ranscripts encoding different isoforms.

基因 ID	968
基因名	CD68
Swiss	P34810
别名	CD68;GP110;LAMP4;SCARD1

产品验证



Western blot analysis of CD68 expressed in Mouse liver,U-87MG,Raji using CD68 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocki ng buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn