

IL7R alpha/CD127 Rabbit mAb

货号: B31328

产品信息

展的 Human,Mouse,Rat Rabbit 文牌性 Monoclonal (機調反应 ・		
克隆性 Monoclonal 預測反应 の用	反应	Human,Mouse,Rat
度用 WB FC 推荐浓度 WB: 1:500 - 1:2000 FC: 1:20 - 1:50 理论分子嚴 65-90kDa 实测分子最 75kDa 形式 Liquid 保存条件 Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. 佩联物 Unconjugated 阳性对照 HepG2,PC-3,Mouse liver,Mouse thymus 细胞定位 Cell membrane,Secreted,Single-pass type I membrane protein,Single-pass type I membrane protein	宿主	Rabbit
应用 WB FC 推荐浓度 WB: 1:500 - 1:2000 FC: 1:20 - 1:50 理论分子量 65-90kDa 实测分子量 75kDa 形式 Liquid 保存条件 Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. 俱联物 Unconjugated 阳性对照 HepG2,PC-3,Mouse liver,Mouse thymus 细胞定位 Cell membrane,Secreted,Single-pass type I membrane protein,Single-pass type I membrane protein	克隆性	Monoclonal
# # # # # # # # # # # # # # # # # # #	预测反应	
FC: 1:20 - 1:50 理论分子量 65-90kDa ***********************************	应用	WB FC
実測分子量 75kDa 形式 Liquid 保存条件 Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. 個联物 Unconjugated 阳性对照 HepG2,PC-3,Mouse liver,Mouse thymus 细胞定位 Cell membrane,Secreted,Single-pass type I membrane protein,Single-pass type I membrane protein	推荐浓度	
形式 Liquid 保存条件 Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. 【 Unconjugated 阳性对照 HepG2,PC-3,Mouse liver,Mouse thymus 如胞定位 Cell membrane,Secreted,Single-pass type I membrane protein,Single-pass type I membrane protein	理论分子量	65-90kDa
保存条件 Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. (实测分子量	75kDa
Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. (周联物 Unconjugated 阳性对照 HepG2,PC-3,Mouse liver,Mouse thymus 如胞定位 Cell membrane,Secreted,Single-pass type I membrane protein,Single-pass type I membrane protein	形式	Liquid
阳性对照 HepG2,PC-3,Mouse liver,Mouse thymus 细胞定位 Cell membrane,Secreted,Single-pass type I membrane protein,Single-pass type I membrane protein	保存条件	
细胞定位 Cell membrane,Secreted,Single-pass type I membrane protein,Single-pass type I membrane protein	偶联物	Unconjugated
	阳性对照	HepG2,PC-3,Mouse liver,Mouse thymus
纯化 Affinity purification	细胞定位	Cell membrane,Secreted,Single-pass type I membrane protein,Single-pass type I membrane protein
	纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human IL7R alpha/CD127.
序列	PESFLDCQIHRVDDIQARDEVEGFLQDTFPQQLEESEKQRLGGDVQSPNCPSEDVVITPESFGRDSSLTCLAGNVSACDA PILSSSRSLDCRESGKNGPHV

靶点信息

研究背景	The protein encoded by this gene is a receptor for interleukin 7 (IL7). The function of this receptor require s the interleukin 2 receptor, gamma chain (IL2RG), which is a common gamma chain shared by the receptors of various cytokines, including interleukins 2, 4, 7, 9, and 15. This protein has been shown to play a critical role in V(D)J recombination during lymphocyte development. Defects in this gene may be associated with severe combined immunodeficiency (SCID). Alternatively spliced transcript variants have been found. [provided by RefSeq, Dec 2015]
------	--

基因 ID	3575
基因名	IL7R
Swiss	P16871
别名	CD127; CDW127; IL-7R-alpha; IL7RA; ILRA

产品验证



Western blot analysis of IL7R alpha/CD127 expressed in Mouse liver, HepG2, PC-3 using IL7R alpha/C D127 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysat es/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn