

Aconitase 2 Rabbit mAb

货号: B31359

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	85kDa
实测分子量	85kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Jurkat,Mouse heart,Rat kidney
细胞定位	Mitochondrion
纯化	Affinity purification
形式 保存条件 偶联物 阳性对照 细胞定位	Liquid Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. Unconjugated HeLa,Jurkat,Mouse heart,Rat kidney Mitochondrion


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Aconitase 2.
序列	IAGTLKFNPETDYLTGTDGKKFRLEAPDADELPKGEFDPGQDTYQHPPKDSSGQHVDVSPTSQRLQLLEPFDKWDGKDL EDLQILIKVKGKCTTDHISAAGPWLKFRGHLDNISNNLLIGAINIENGKANSVRNAVTQEFGPVPDTARYYKKHGIRWVVIG DENYGEGSSREHAALEPRHLGGRAIITKSFARIHETNLKKQGLLPLTFADPADYNKIHPVDKLTIQGLKDFTPGKPLKCIIKHP NGTQETILLNHTFNETQIEWFRAGSALNRMKELQQ

靶点信息

研究背景	The protein encoded by this gene belongs to the aconitase/IPM isomerase family. It is an enzyme that cat alyzes the interconversion of citrate to isocitrate via cis-aconitate in the second step of the TCA cycle. This protein is encoded in the nucleus and functions in the mitochondrion. It was found to be one of the mitochondrial matrix proteins that are preferentially degraded by the serine protease 15(PRSS15), also known as Lon protease, after oxidative modification.
基因 ID	50
基因名	ACO2
Swiss	Q99798
别名	ACO2;ACONM;HEL-S-284;ICRD;OCA8;OPA9

产品验证



Western blot analysis of Aconitase 2 expressed in Mouse heart,Rat kidney,HeLa using Aconitase 2 Rab bit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure ti me: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn