

MTH1 Rabbit mAb

货号: B31372

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	17kDa/19kDa/20kDa/22kDa
实测分子量	18kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SW480,K562,22RV1,THP-1,HepG2,SKOV3,293T
细胞定位	Cytoplasm,Mitochondrion matrix,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human MTH1.
序列	MSGISPQQMGEPEGWSGKNPGTMGASRLYTLVLVLQPQRVLLGMKKRGFGAGRWNFGGKVVQEGETIEDGARRELQ EESGLTVDALHKVGQIVFEFVGEPELMDVHVFCDSIQGTPVESDEMPCWFQLDQIPFKDMWPDDSYWFPLLLQKKKF HGYFKFQGQDTILDYTLREVDTV

靶点信息

研究背景	Misincorporation of oxidized nucleoside triphosphates into DNA/RNA during replication and transcription can cause mutations that may result in carcinogenesis or neurodegeneration. The protein encoded by this gene is an enzyme that hydrolyzes oxidized purine nucleoside triphosphates, such as 8-oxo-dGTP, 8-oxo-dATP, 2-hydroxy-dATP, and 2-hydroxy rATP, to monophosphates, thereby preventing misincorporation. The encoded protein is localized mainly in the cytoplasm, with some in the mitochondria, suggesting that it is involved in the sanitization of nucleotide pools both for nuclear and mitochondrial genomes. Several alternatively spliced transcript variants, some of which encode distinct isoforms, have been identified. Additional variants have been observed, but their full-length natures have not been determined. A single-nucleotide polymorphism that results in the production of an additional, longer isoform (p26) has been described.
基因ID	4521
基因名	NUDT1
Swiss	P36639
别名	NUDT1;MTH1

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn