

Wnt2b Rabbit mAb

货号: B31380

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	44kDa
实测分子量	44kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,PC-3,U-87MG,Mouse lung,Mouse testis,Mouse brain,Rat testis,Rat uterus
细胞定位	extracellular region,extracellular space
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Wnt2b.
序列	KAFVDAKEKRLKDARALMNLHNNRCGRTAVRRFLKLECKCHGVSGSCTLRTCWRALSDFRRTGDYLRYYDGAVQVMA TQDGANFTAARQGYRRATRTDLV

靶点信息

研究背景	This gene encodes a member of the wingless-type MMTV integration site (WNT) family of highly conserved, secreted signaling factors. WNT family members function in a variety of developmental processes including regulation of cell growth and differentiation and are characterized by a WNT-core domain. This gene may play a role in human development as well as carcinogenesis. Alternative splicing results in multiple transcript variants. [provided by RefSeq, May 2014]
------	--

基因ID	7482
基因名	WNT2B
Swiss	Q93097
别名	WNT13

产品验证



Western blot analysis of Wnt2b expressed in Mouse lung, Rat testis, 293T using Wnt2b Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn