

Phospho-AKT1 (S129) Rabbit mAb

货号: B31389

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	60kDa
实测分子量	60kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	Cell membrane,Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Phospho-AKT1 (S129).
序列	DNSGA


靶点信息

研究背景

The serine-threonine protein kinase encoded by the AKT1 gene is catalytically inactive in serum-starved p rimary and immortalized fibroblasts. AKT1 and the related AKT2 are activated by platelet-derived growth factor. The activation is rapid and specific, and it is abrogated by mutations in the pleckstrin homology do main of AKT1. It was shown that the activation occurs through phosphatidylinositol 3-kinase. In the devel oping nervous system AKT is a critical mediator of growth factor-induced neuronal survival. Survival factors can suppress apoptosis in a transcription-independent manner by activating the serine/threonine kinase AKT1, which then phosphorylates and inactivates components of the apoptotic machinery. Mutations in this gene have been associated with the Proteus syndrome. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jul 2011]

基因ID	207
基因名	AKT1
Swiss	P31749
别名	AKT; CWS6; PKB; PKB-ALPHA; PRKBA; RAC; RAC-ALPHA

产品验证



Western blot analysis of Phospho-AKT1 (S129) expressed in 293T using Phospho-AKT1 (S129) Rabbit m Ab at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 12 0s.

实验步骤

访问官网浏览详情: www.ablybio.cn