

# Phospho-Bad (S112) Rabbit mAb

货号: B31392

# 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IP
推荐浓度	WB: 1:500 - 1:2000 IP: 1:20 - 1:50
理论分子量	18kDa
实测分子量	23kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,A-549,Mouse brain,Mouse thymus,Mouse lung,Rat brain
细胞定位	Cytoplasm,Mitochondrion outer membrane
纯化	Affinity purification

# 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Phospho-Bad (S112).
序列	MFQIPEFEPSEQEDSSSAERGLGPSPAGDGPSGSGKHHRQAPGLLWDASHQQEQPTSSSHHGGAGAVEIRSRHSSYPA GTEDDEGMGEEPSPFRGRSRSA

#### 靶点信息

研究背景	The protein encoded by this gene is a member of the BCL-2 family. BCL-2 family members are known to be regulators of programmed cell death. This protein positively regulates cell apoptosis by forming heterod imers with BCL-xL and BCL-2, and reversing their death repressor activity. Proapoptotic activity of this protein is regulated through its phosphorylation. Protein kinases AKT and MAP kinase, as well as protein phosphatase calcineurin were found to be involved in the regulation of this protein. Alternative splicing of this gene results in two transcript variants which encode the same isoform. [provided by RefSeq, Jul 2008]
------	---

基因ID	572
基因名	BAD
Swiss	Q92934
别名	BAD;BBC2;BCL2L8;Bad

# 产品验证



Western blot analysis of Phospho-Bad (S112) expressed in Mouse brain,Rat brain,HeLa using Phosp ho-Bad (S112) Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:50 00. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL E nhanced Kit. Exposure time: 120s.

### 实验步骤

访问官网浏览详情: www.ablybio.cn