

CD105 Rabbit mAb

货号: B31409

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	90kDa
实测分子量	90kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HUVEC
细胞定位	Membrane,Single-pass type I membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human CD105.
序列	MDRGTLPLAVALLLASCSLSPTSLAETVHCDLQPVGPERGEVTYTTSQVSKGCVAQAPNAILEVHVLFLEFPTGPSQLELTL QASKQNGTWPREVLLVLSVNSSVFLHLQALGIPLHLAYNSSLVTFQEPPGVNTTELPSFPKTQILEWAAERGPITSAAELND PQSILLRLGQAQGSLSFCMLEASQDMGRTLEWRPRT

靶点信息

研究背景	This gene encodes a homodimeric transmembrane protein which is a major glycoprotein of the vascular e ndothelium. This protein is a component of the transforming growth factor beta receptor complex and it b inds to the beta1 and beta3 peptides with high affinity. Mutations in this gene cause hereditary hemorrha gic telangiectasia, also known as Osler-Rendu-Weber syndrome 1, an autosomal dominant multisystemic vascular dysplasia. This gene may also be involved in preeclampsia and several types of cancer. Alternati vely spliced transcript variants encoding different isoforms have been found for this gene. [provided by R efSeq, May 2013]
基因 ID	2022
基因名	ENG
Swiss	P17813
别名	END;HHT1;ORW1;CD105;ENG;endoglin

产品验证



Western blot analysis of CD105 expressed in HUVEC using CD105 Rabbit mAb at 1:1000. Secondary antib ody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn