

Adenylate kinase 4 Rabbit mAb

货号: B31441

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	25kDa
实测分子量	25kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,HeLa,SH-SY5Y
细胞定位	Mitochondrion matrix
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Adenylate kinase 4.
序列	MASKLLRAVILGPPGSGKGTVCQRIAQNFGLQHLSSGHFLRENIKASTEVGEMAKQYIEKSLLVPDHVITRLMMSELENRR GQHWLLDGFPRTLGQAEAL

靶点信息

研究背景	This gene encodes a member of the adenylate kinase family of enzymes. The encoded protein is localized to the mitochondrial matrix. Adenylate kinases regulate the adenine and guanine nucleotide composition s within a cell by catalyzing the reversible transfer of phosphate group among these nucleotides. Five iso zymes of adenylate kinase have been identified in vertebrates. Expression of these isozymes is tissue-spe cific and developmentally regulated. A pseudogene for this gene has been located on chromosome 17. Th ree transcript variants encoding the same protein have been identified for this gene. Sequence alignment suggests that the gene defined by NM_013410, NM_203464, and NM_001005353 is located on chromoso me 1.
基因ID	205
基因名	AK4
Swiss	P27144
别名	AK4; AK 4; AK3; AK3L1; AK3L2; adenylate kinase 4

产品验证



Western blot analysis of Adenylate kinase 4 expressed in 293T,HeLa,SH-SY5Y using Adenylate kina se 4 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysate s/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced K it. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn